


Instructions to follow when taking the COVID - 19 vaccine



1

Initiate a rapid response and approach the vaccination centers as soon as you are notified that you are within the target group for the vaccine.



2

Commitment to taking the second dose of the vaccine after the specified period from the date of taking the first dose, in order to obtain the desired effectiveness, as one dose of the vaccine is not sufficient.



3

Be aware that the effect of the vaccine begins 14 days after taking the second dose to get the desired effect, and before this period the person is still susceptible to infection and he should go to the health institution if he have any suspected symptoms.

Attention


4

To reduce the side effects of vaccines:

Use fever-reducing medications such as paracetamol (Adol) to reduce fever, headache and joint pain every 8-6 hours if needed.



Apply cold compresses on the injection site to reduce pain, redness and swelling at the injection site, if any.



If these symptoms persist or if you experience severe side effects, please go to the nearest health institution.


5

After taking the vaccine doses, the person remains in the vaccination centers for 15 minutes to observe and to make sure that no dangerous side effect occurs, which requires direct medical intervention such as an attack of hypersensitivity.



6

Please take care for Continuing to adhere to other preventive measures in place, such as washing hands, wearing a mask, and physical distancing, after taking the vaccine doses, as taking the vaccine does not nullify these procedures.



Not to heed rumors and take all the information about the vaccine from reliable official sources.